

Table 1

Incyte Project ID	Polypeptide		Incyte		Polynucleotide		Incyte
	SEQ ID NO:	SEQ ID NO:	Polypeptide ID	Polypeptide ID	SEQ ID NO:	SEQ ID NO:	Polynucleotide ID
2515666	1		2515666CD1		2		2515666CB1

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability Score	GenBank Homolog
1	251566CD1	g181382	6.90E-270	Cytochrome P450 [Homo sapiens] (Ikeya, K. et al. (1989) Mol. Endocrinol. 3:1399-1408)

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	2515666CD1	504	S10 S89 S212 T267 T323 T378 S69 T93 S279 S348 T363 S433 T486		E-class P450 group I PR00463: S69-L68, T93-S114, S195-S213, K294-T311, I314-G340, E357-P375, N398-T422, M436-C446, C446-F469 E-class P450 group II PR00464: G130-A150, K294-V322, T323-G340, Q352-F372, K392-E407, L408-A423, S433-C446, C446-F469 P450 superfamily signature PR00385: A305-V322, T323-D336, A358-F369, M437-C446, C446-F457 Cytochrome P450: F439-G448 CYTOCHROME P450 DM00022 S43414 18-491: V24-L485 Cytochrome P450 PD000021: L111-R284, F263-G386, F387-P469 Cytochrome P450: P42-L499 Cytochrome P450 cysteine heme-iron ligand signature: F415-E468 Cytochrome P450 cysteine heme-iron ligand signature BL000086: M436-L467 Transmembrane domain: L14-R32 Signal peptide: M1-C22	BLIMPS- PRINTS BLIMPS- PRINTS BLIMPS- PRINTS BLIMPS- PRINTS BLAST-DOMO BLAST- PRODOM HMMER-PFAM PROFILES- SCAN BLIMPS- BLOCKS HMMER HMMER

TABLE 20 "022303000

Table 4

Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID	Sequence Length	Selected Fragments	Sequence Fragments	5' Position	3' Position
2	2515666CBI	1790	1-77, 891-915	SZAE00257F1	1356	1790
				SZAE00059F1	142	770
				2515666H1 (LIVRTUT04)	1	177
				SZAE00073F1	905	1601
				SZAE00189F1	251	871
				SZAE00140F1	843	1582

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
2	251566GB1	LIVRTUT04

Table 6

Library	Vector	Library Description
LIVRT04	pINCY	Library was constructed using RNA isolated from liver tumor tissue removed from a 50-year-old Caucasian male during a partial hepatectomy. Pathology indicated a grade 3-4 hepatoma, forming a mass. Patient history included benign hypertension and hepatitis. Hepatitis B core antigen and hepatitis B surface antigen was present in the patient.

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and tsearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fasta E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322; Durbin, R. et al. (1998) Our World View, in a Nussli, Cambridge Univ. Press, pp. 1-350.	PFAM hits: Probability value= 1.0E-3 or less Signal peptide hits: Score= 0 or greater

Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribkov, M. et al. (1988) CABIOS 4:61-66; Gribkov, M. et al. (1989) Methods Enzymol. 183:146-159; Baroch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score>GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phis Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Baroch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59. Genetics Computer Group, Madison, WI.	